



An evidence for surface expression of an immunogenic epitope of sarcoplasmic/endoplasmic reticulum calcium-ATPase2a on antigen-presenting cells from naive mice in the mediation of autoimmune myocarditis



Rajkumar Arumugam^a, Bharathi Yalaka^{a,1}, Chandirasegaran Massilamany^{a,2},
M.S. Shihabudeen Haider Ali^b, Ninaad Lasrado^a, Sabarirajan Jayaraja^a, Jean-Jack Riethoven^c,
Xinghui Sun^b, Jay Reddy^{a,*}

^a School of Veterinary Medicine and Biomedical Sciences, University of Nebraska-Lincoln, Lincoln, NE, United States

^b Department of Biochemistry, University of Nebraska-Lincoln, Lincoln, NE, United States

^c Center for Biotechnology, University of Nebraska-Lincoln, Lincoln, NE, United States

ARTICLE INFO

Keywords:

Autoreactive T cells
Self-antigens
Myocarditis
Autoimmunity
Mouse model
APCs
T cells and SERCA2a

ABSTRACT

We recently reported identification of sarcoplasmic/endoplasmic reticulum calcium-ATPase2a (SERCA2a) 971-990, which induces atrial myocarditis by generating autoreactive T cells in A/J mice. However, it was unknown how antigen-sensitized T cells could recognize SERCA2a 971-990, since SERCA2a-expression is confined to an intracellular compartment. In this report, we present evidence that antigen-presenting cells (APCs) from lymphoid and non-lymphoid organs in naïve animals present SERCA2a 971-990 and stimulate antigen-specific T cells. Using major histocompatibility complex (MHC) class II dextramers for SERCA2a 971-990, we created a panel of T cell hybridomas and demonstrated that splenocytes from naïve A/J mice stimulated the hybridoma cells without exogenous supplementation of SERCA2a 971-990. We then recapitulated this phenomenon by using SERCA2a 971-990 -specific primary T cells, verifying that the T cell responses were MHC-restricted. Furthermore, SERCA2a 971-990 -sensitized T cells exposed to APCs from naïve mice were found to produce the inflammatory cytokines interferon- γ , granulocyte macrophage colony stimulating factor, and interleukin-17A, which are implicated in the induction of myocarditis. Finally, while T cells exposed to mononuclear cells (MNCs) obtained from heart and liver also responded similarly to splenocytes, endothelial cells (ECs) generated from the corresponding organs displayed opposing effects, in that the proliferative responses were suppressed with the heart ECs, but not with the liver ECs. Taken together, our data suggest that the surface expression of SERCA2a 971-990 by naïve APCs can potentially trigger pathogenic autoreactive T cell responses under conditions of autoimmunity, which may have implications in endothelial dysfunction.

Abbreviations: SERCA2a, sarcoplasmic/endoplasmic reticulum Ca²⁺-ATPase2a; MHC, major histocompatibility complex; APCs, antigen presenting cells; ECs, endothelial cells; RNase, bovine ribonuclease; MCC, moth cytochrome C; FBS, fetal bovine serum; ³[H], tritiated-thymidine; cpm, counts per minute; IL, interleukin; SA, streptavidin; 7-AAD, 7-aminoactinomycin D; RT, room temperature; TCR, T cell receptor; PBS, phosphate-buffered saline; MNCs, mononuclear cells; IFN, interferon; TNF, tumor necrosis factor; GM-CSF, granulocyte macrophage colony stimulating factor; SR, sarcoplasmic reticulum; Th, T helper; PI, propidium iodide; PD-1, programmed cell death protein 1; PD-L1, programmed death-ligand 1; PMA, phorbol 12-myristate 13-acetate; PLP, proteolipid protein; MBP, myelin basic protein; DNase1, deoxyribonuclease I; PECAM, platelet endothelial cell adhesion molecule; Sf, *Spodoptera frugiperda*; DMEM, dulbecco's modified eagle medium; HBSS, hank's balanced salt solution; CCR, chemokine receptor; LFA, lymphocyte function associated antigen; FITC, fluorescein isothiocyanate; APC, allophycocyanin; PE, phycoerythrin

* Corresponding author at: Room 202, Bldg. VBS, University of Nebraska-Lincoln, Lincoln, NE 68583, United States.

E-mail address: nreddy2@unl.edu (J. Reddy).

¹ Present address: Bristol-Myers Squibb – Hopewell, Pennington, NJ.

² Present address: CRISPR Therapeutics, Cambridge, MA.

<https://doi.org/10.1016/j.imbio.2019.12.005>

Received 25 August 2019; Received in revised form 6 December 2019; Accepted 7 December 2019

Available online 13 December 2019

0171-2985/ © 2019 Elsevier GmbH. All rights reserved.

1. Introduction

Autoimmune diseases can result from autoreactive T cells or B cells or both, but detection of autoreactive cells in healthy individuals does not mean that clinical manifestations would ensue in their lifetime. It is well established that developing T and B lymphocytes are expected to recognize self-antigens during maturation processes in their generative organs, thymus and bone marrow, respectively, prior to their export to the periphery. Because weakly recognized lymphocytes are allowed to exit the primary organs as a result of positive selection, detection of autoreactive cells in the periphery would not be surprising (Theofilopoulos et al., 2017). Yet, most individuals remain healthy, and how such a tolerance can be maintained is a fundamental question in autoimmunity research.

Mechanistically, various theories have been proposed to suggest that self-tolerance can be broken in genetically susceptible individuals under altered environmental conditions, which may include both exogenous (e.g., exposure to microbial infections) and endogenous (e.g., defects in the maturation processes, faulty regulation) factors (Theofilopoulos et al., 2017; Cheng and Anderson, 2018). It is widely believed that the frequencies of antigen-specific lymphocytes are in the range of 1 in 1×10^5 to 1×10^6 cells in healthy individuals (Goodnow et al., 2005). Despite such low frequencies, upon exposure to microbial infections, lymphocytes faithfully respond to foreign antigens, which is one of the cardinal features of adaptive immune cells. By contrast, autoreactive lymphocytes, although present in similar frequencies, are not expected to react to self-antigens (Lohse et al., 1996; Alanio et al., 2010; Steinert et al., 2012; Vrisekoop et al., 2017), even though they continuously see the antigens. Whether this continuous exposure is a critical requirement for self-reactive lymphocytes to become tolerant in the periphery is not clear. In support of this proposition, however, transgenic expression of self-antigens has been shown to promote tolerance (Skowronski et al., 1990; French et al., 1997; Bridgett et al., 1998). Conversely, it has been held that either autoreactive cells recognizing self-antigens undergo apoptosis or such recognition can be suppressed by the mediation of regulatory T cells (Theofilopoulos et al., 2017; Sakaguchi et al., 1995).

Experimentally, use of animal models has enhanced our understanding of how self-tolerance can be broken, leading to the induction of pathogenic responses. We have been engaged in determining the role of antigen-specific T cells in the causation of organ-specific diseases, in particular, heart (Krishnan et al., 2017; Krishnan et al., 2018; Basavalingappa et al., 2016; Basavalingappa et al., 2017; Gangaplara et al., 2012). This work led us to identify at least three intracellular proteins – adenine nucleotide translocator (Basavalingappa et al., 2016), branched chain α -ketoacid dehydrogenase kinase (Krishnan et al., 2017), and sarcoplasmic/endoplasmic reticulum calcium-ATPase2a (SERCA2a) (Krishnan et al., 2018) – as autoimmune targets in the development of myocarditis/dilated cardiomyopathy in myocarditis-susceptible, A/J mice. SERCA2a is unique in that it contains at least six immunodominant T cell epitopes. One of these, SERCA2a 971-990, can bind two major histocompatibility complex (MHC) class II alleles (IA^k and IE^k) with varied affinities and induce mainly atrial myocarditis by generating antigen-specific T cells that can transfer disease to naïve animals (Krishnan et al., 2018). While preferential induction of myocardial lesions in the atria could be correlated with enhanced expression of SERCA2a in the atria rather than the ventricles (Krishnan et al., 2018), it was unknown whether SERCA2a peptide can be displayed by the naïve antigen-presenting cells (APCs) to trigger autoreactive T cell responses. To address this question, we generated SERCA2a 971-990 -specific T cell hybridomas and demonstrated that hybridoma cells respond to SERCA2a 971-990 when stimulated by splenocytes from naïve mice without exogenous supplementation of peptide. While we confirmed these findings with primary T cells specific to SERCA2a 971-990, we unexpectedly noted that the endothelial cells (ECs) derived from hearts and livers from naïve mice modulated

SERCA2a 971-990 -reactive T cell responses differentially, which may have implications in our understanding of endothelial dysfunction that might occur in vascular inflammation and coronary heart diseases.

2. Materials and methods

2.1. Mice

Male A/J mice (6- to 8-wk-old, H-2^b) obtained from the Jackson Laboratory (Bar Harbor, ME) were maintained in accordance with the Institutional Animal Care and Use Committee guidelines, University of Nebraska-Lincoln, Lincoln, NE. Euthanasia was performed using carbon dioxide as recommended by the Panel on Euthanasia, American Veterinary Medical Association.

2.2. Peptide synthesis

SERCA2a 971-990 (Ac-KISLPVILMDETLKFVARNY), bovine ribonuclease (RNase) 43-56 (VNTFVHESLADVQA), and moth cytochrome C (MCC) 82-103 (FAGLKKANERADLIAYLKQATK) (GenScript, Piscataway, NJ) were synthesized by 9-fluorenylmethoxycarbonyl chemistry. The purity of peptides was more than 90 % as evaluated by high-performance liquid chromatography, and their identities were confirmed by mass spectroscopy. Peptides dissolved in ultra-pure water were aliquoted and stored at -20°C until further use.

2.3. Immunization procedures, and generation of primary T cell cultures

On days 0 and 7, animals were immunized subcutaneously in the inguinal and sternum regions with emulsions of SERCA2a 971-990 (50 $\mu\text{g}/\text{animal}$) prepared in complete Freund's adjuvant containing *Mycobacterium tuberculosis* H37RA extract (5 mg/ml; Difco Laboratories, Detroit, MI, USA) (Cihakova et al., 2008; Massilamany et al., 2016). At termination on day 21 post-immunization, single cell suspensions were prepared from the draining lymph nodes. Cells were stimulated with SERCA2a 971-990 (20 $\mu\text{g}/\text{ml}$) at a density of 5×10^6 cells/ml for 2 days in growth medium (RPMI medium supplemented with 10 % fetal bovine serum [FBS], 1 mM sodium pyruvate, 4 mM L-glutamine, 1x each of non-essential amino acids and vitamin mixture, and 100 U/ml penicillin-streptomycin [Lonza, Walkersville, MD, USA]) (Massilamany et al., 2016), and growth medium containing interleukin (IL)-2 was then added. Cells were then stimulated 3–4 times with SERCA2a 971-990 as above, using syngeneic APCs.

2.4. Creation of MHC class II/ IA^k or IE^k dextramers and dextramer staining

We recently reported the generation of MHC dextramers, more sensitive reagents than tetramers, and used them to detect antigen-specific T cells in a variety of experimental systems (Krishnan et al., 2017; Krishnan et al., 2018; Basavalingappa et al., 2017; Massilamany et al., 2015; Massilamany et al., 2011a; Massilamany et al., 2011b; Gangaplara et al., 2013). We also had demonstrated their utility in generating T cell hybridomas for the central nervous system antigen proteolipid protein (PLP) 139-151 (Krishnan et al., 2015). We adopted a similar approach to create SERCA2a 971-990 -specific T cell hybridomas. Essentially, A/J mice express two MHC class II alleles: IA^k and IE^k (Basavalingappa et al., 2017). We generated two sets of MHC class II dextramers, with one set each for IA^k (SERCA2a 971-990 /RNase 43-56) and IE^k (SERCA2a 971-990 /MCC 82-103), where RNase 43-56 and MCC 82-103 were used as controls for IA^k and IE^k molecules, respectively (Krishnan et al., 2018). The sequences for all the above were covalently tethered to the β chains of the corresponding MHC class II alleles. In brief, α and β constructs of IA^k and IE^k molecules were expressed in baculovirus in Sf9 insect cells. After affinity-column purifications using anti- IA^k and anti- IE^k antibodies, soluble IA^k /SERCA2a 971-990 and RNase 43-56, as well as IE^k /SERCA2a 971-990 and MCC

82-103 monomers, were biotinylated, and dextramers were derived using streptavidin (SA)/fluorophore-conjugated dextran molecules as we have described previously (Massilamany et al., 2011b). For dextramer staining, cells were stained with IA^k- (SERCA2a 971-990 and RNase 43-56) or IE^k- (SERCA2a 971-990 and MCC 82-103) dextramers, followed by anti-CD4 FITC (Fluorescein isothiocyanate) and 7-aminocoumarin-6 (7-AAD). After washing, cells were acquired by flow cytometry, and the percent dextramer⁺ cells were analyzed using FlowJo software (Tree Star, Ashland, OR, USA) (Massilamany et al., 2011b; Massilamany et al., 2011c; Reddy et al., 2003).

2.5. Derivation of T cell hybridomas specific to SERCA2a 971-990

By using MHC class II dextramers as screening and sorting tools, we generated a panel of 92 T cell hybridoma clones as we have described previously (Krishnan et al., 2015), and we selected one clone (80-P8-8-E2), designated hereafter, clone 80 for further characterization. While, antigen-specificity of this clone was determined based on dextramer staining as described above (Krishnan et al., 2015), expression of various markers was analyzed by standard flow cytometry.

2.6. Intracellular cytokine staining

T cell hybridomas were briefly stimulated for 4–5 hours with an activation cocktail containing phorbol 12-myristate 13-acetate (PMA), ionomycin, and Brefeldin A (Biolegend). After staining with anti-CD4 FITC and 7-AAD, cells were fixed and permeabilized, followed by staining with cytokine antibodies as recommended by the manufacturer (Biolegend). The following cytokine antibodies used were (clone numbers indicated in parentheses): IL-2 (JES6); interferon (IFN)- γ (XMG1.2) (Th1 subset); IL-4 (11B11); IL-5 (TRFK5); IL-13 (eBio13A) (Th2 subset); IL-17A (eBio17B7); IL-17 F (eBio18F10); IL-22 (IH8PWSR) (Th17 subset); IL-6 (MP5-20F3); tumor necrosis factor (TNF)- α (MP6-X722); and granulocyte macrophage colony stimulating factor (GM-CSF) (MP1-22E9) (other inflammatory cytokines). In experiments involving T cells sensitized with SERCA2a 971-990, T cells were cultured with splenocytes from naïve mice in the presence or absence of SERCA2a 971-990 for two days where monensin (1x) was added in the last 12 h at each of the four time points namely, 6, 12, 24 and 36 h. After washing and staining with anti-CD4-FITC and 7-AAD, cells were fixed and permeabilized and stained with three cocktails of cytokine antibodies namely, IFN- γ -allophycocyanin (APC) and IL-2-phycoerythrin (PE), IL-17A-APC and GM-CSF-PE and their corresponding controls (rat IgG1-APC and rat IgG2a/IgG2b-PE). After acquiring the cells by flow cytometry, percentages of cytokine-producing, viable (7-AAD⁻) cells within the CD4 subset were analyzed using FlowJo software. In all the treatment groups, while, APC-conjugated, IFN- γ , IL-17A and isotype controls are shown in the Y axis, the PE-conjugated, IL-2, GM-CSF and isotype controls are shown in the X axis in the flow cytometric plots. To determine the frequencies of IFN- γ , IL-17A and their controls, percentage values of the upper left and upper right quadrants were added in their corresponding flow cytometric plots. Similarly, percentage values in the upper right and lower right quadrants were added to obtain the frequencies of cells producing IL-2, GM-CSF and their controls, in their respective plots.

2.7. Proliferative responses of SERCA2a 971-990 -specific T cell hybridoma and primary T cells to splenocytes from naïve mice

To determine whether APCs spontaneously stimulate SERCA2a 971-990 -specific T cells, we used both hybridoma cells (clone 80) and primary T cells. For hybridomas, cells were cultured for 24 h (2.5×10^4 cells/ml) with splenocytes obtained from naïve A/J mice at different ratios (1:1 to 1:20) in the presence or absence of SERCA2a 971-990 or control (MCC 82-103) (50 μ g/ml). After pulsing with ³[H] thymidine for 16 h, proliferative responses were measured as cpm. Similarly,

primary T cells were also cultured with splenocytes at a ratio of 1:1 with or without peptides for 2 days, and proliferative responses were measured 16 h after pulsing with ³[H] thymidine (Rock et al., 1990; Smith and Allen, 1992; Donermeyer et al., 1995). In MHC blocking experiments, Fc-blocker-treated splenocytes from naïve mice were cultured with or without T cells supplemented with anti-IA^k (10-2.16, clone TIB 93, 20 μ g/ml) and anti-IE^k (M5/114.15.2, clone TIB 120, 100 μ g/ml; American Tissue Culture Collection, Manassas, VA, USA) or Rat IgG (control; 100 μ g/ml, MP Biomedicals, Santa Ana, CA, USA). The proliferative responses were measured as above (Tamura et al., 1996; Bright et al., 1999).

2.8. Determination of SERCA2a 971-990 -reactive T cell responses to mononuclear cells (MNCs) generated from heart and liver

Groups of 5 to 6 naïve A/J mice were euthanized, and animals were immediately perfused with 10 ml of ice-cold 1x phosphate-buffered saline (PBS) prior to harvesting of hearts and livers (Krishnan et al., 2017). Briefly, hearts were cut into two halves and the pooled tissues were minced and transferred to gentleMACS C Tube (Miltenyi Biotec, San Diego, CA, USA) containing 4.7 ml of Hank's Balanced Salt Solution supplemented with Collagenase II (600 U/ml; Worthington Biochemical Corporation, Lakewood, NJ, USA) and DNase I (60 U/ml; AppliChem GmbH, Darmstadt, Germany). After mixing by MACS Dissociator, samples were incubated for 30 min at 37 °C using the MACSmix Tube Rotator (Miltenyi Biotec); the tissue suspensions were mixed using the MACS Dissociator. The homogenate was centrifuged and cell suspensions were filtered using a 70- μ m cell strainer; after washing, red blood cells were lysed and the debris removed using Debris Removal Solution as recommended (Miltenyi Biotec). Finally, cell pellets were re-suspended in growth medium, and cells were treated with Fc blocker (Biolegend), followed by staining with 7-AAD and anti-CD45. After washing, CD45⁺ cells within the live subset (7-AAD⁻) were sorted by Aria flow cytometry (FACS Aria, BD Biosciences, San Jose, CA, USA). To obtain single cell suspensions from livers, tissues were minced and centrifuged. The homogenate was digested in Hank's Balanced Salt Solution containing Collagenase II (Worthington Biochemical Corporation) and DNase I (AppliChem GmbH) at 37 °C for 30 min, followed by centrifugation to remove hepatocytes. Cell pellets were washed with 1x PBS, and cell suspensions were then subjected to Percoll density gradient centrifugation. The interphase representing the MNCs was harvested, and, after washing and treating with red blood cell lysis buffer, samples were centrifuged to obtain the MNC fraction. Cells were then treated with Fc blocker, and after staining with CD45 and 7-AAD, CD45⁺ cells were sorted by flow cytometry as above. Finally, cells derived from both hearts and livers were evaluated for CD11b, CD11c, CD19, and IE^k, as well as for T cell responses, using SERCA2a 971-990 -specific primary T cells by ³[H] thymidine-incorporation assay as described above.

2.9. Isolation of primary ECs from hearts and livers and evaluation of T cell responses to the ECs

ECs from hearts and livers of groups of 5–6 naïve A/J mice were isolated using the previously published method with some modifications (Lim et al., 2003; Chou et al., 2015; van Beijnum et al., 2008). In brief, animals were euthanized using 30 % isoflurane, and ECs were isolated by positive selection using immunomagnetic beads specific to PECAM-1 (CD31) (Daneker et al., 1998; McDouall et al., 1996; Grafe et al., 1994; Grafe et al., 1993). Essentially, the CD31-bound magnetic beads were freshly prepared a day prior to isolating the cells by mixing sheep anti-rat IgG Dynabeads (Thermo Fisher Scientific) and anti-CD31 (Clone: MEC13.3, Biolegend) in a final volume of 500 μ l after a series of washing steps. Finally, the CD31-conjugated beads were washed 4 times and resuspended in a volume of 200 μ l for further use. To isolate ECs, heart and liver tissues were minced with scissors to make small

pieces (1 to 2 mm³) and digested with Collagenase II and Dispase (1 mg/ml each in DMEM/F12; GE Healthcare Bio-Sciences AB, Uppsala, Sweden) at 37 °C for 40 min. The digested tissues were washed with DMEM/F12 medium containing 10 % FBS; suspensions were filtered using 70- μ m and 40- μ m meshes and centrifuged at 500 g for 10 min at 4 °C. Cell suspensions were then incubated with the Dynabeads coated with PECAM-1 (CD31) antibody. The bead-bound cells were collected using BD IMag cell separation magnet (BD Biosciences) and washed 3 times. Cells that bound to the magnetic beads were re-suspended and cultured in 0.1 % gelatin-coated tissue culture plates in EC growth medium (Cell Biologics). After growing for 5–7 days, cells were harvested at > 80 % confluence with trypsin-EDTA and used for experiments; purity of ECs was ascertained by anti-CD31 staining. For proliferation experiments, ECs and SERCA2a 971-990 -sensitized T cells were cultured at a ratio of 1:1 individually or together with or without SERCA2a 971-990 or MCC 82-103 for 2 days, and proliferative responses were measured by pulsing the cells with ³[H] thymidine 16 h later.

2.10. IFN- γ treatment of ECs

The CD31⁺ heart and liver ECs plated in gelatin-coated culture flasks (T25 cm²) were grown to ~80 % confluency. On day 7 post-culturing, cells were harvested by trypsinization (0.05 %) and treated with or without Mouse IFN- γ (500 ng/ml) (Biolegend) (Huynh et al., 1995). After 3 days, cells were stained with antibodies for IA^k, IE^k, CD80, CD86, and CD40 molecules and 7-AAD. Cells were acquired by flow cytometry, and percentages of cells positive for each marker were analyzed in the live (7-AAD⁻) subset using FlowJo software.

2.11. Statistics

We used Wilcoxon rank-sum followed by Bonferonni multiple test correction to determine differences in T cell responses. Observations for the T cell responses were normalized prior to statistical testing per repeat experiment based on the average sum of input hybridoma or primary T cells and APCs or ECs. $p \leq 0.05$ was considered significant.

3. Results and discussion

We recently reported that SERCA2a contains multiple antigenic determinants, and one of these epitopes, SERCA2a 971-990, was found

to induce atrial myocarditis in A/J mice by generating both T cell and antibody responses (Krishnan et al., 2018). However, it was unknown as to how SERCA2a can be recognized by autoreactive T cells. Although our efforts to localize the expression of SERCA2a in cardiac myocytes provided leads to predict that its expression may occur close to the plasma membrane (Krishnan et al., 2018), supporting evidence was lacking. Thus, we made an effort to derive T cell hybridomas for SERCA2a 971-990 with the expectation that T cells can respond to antigens if they are displayed by APCs in naïve animals. From a panel of 92 hybridoma clones, we chose clone 80 that was positive for CD3, CD4, TCR $\nu\beta 10b$, CD45, CD5, CD62 L, CD44, CD28, PD-1, PD-L1, CD2, LFA-2 and CCR4. We determined antigen specificity of this clone by staining with IE^k and IA^k dextrans for SERCA2a 971-990, expecting that the IE^k-specific clone 80 hybridoma cells should bind only to IE^k, but not IA^k dextrans or controls (MCC 82-103 or RNase 43-56), and it was the case (Fig. 1A, top panel). We also verified the cytokine-producing ability of hybridoma cells to be in the order of TNF- α , IL-2, IL-4, and IL-6 (Fig. 1B, bottom panel). Similar profiles have been described for other antigens (Krishnan et al., 2015).

3.1. Splenocytes from naïve A/J mice trigger antigen-specific T cell responses spontaneously

We took advantage of SERCA2a 971-990 -specific T cell hybridomas to test a hypothesis that APCs from naïve animals display MHC/SERCA2a complexes that can trigger antigen-specific T cell responses. First, we cultured the clone 80 cells with splenocytes generated from naïve A/J mice in the presence or absence of SERCA2a 971-990 or MCC 82-103 (control) (Ashwell et al., 1988; Merrick et al., 2005; Prasad and Gregerson, 1997). The clone 80 cells responded spontaneously to splenocytes at ratios ranging from 1:1 to 1:20 with ratios up to 1:10 yielded significant differences between treatments (Fig. 2). Responses were antigen specific, since SERCA2a 971-990, but not control (MCC 82-103), stimulated clone 80 cells. Of note, proliferative responses obtained by co-culturing splenocytes with clone 80 cells alone, and with control (MCC 82-103) were also comparable. The data suggest that naïve APCs express SERCA2a 971-990 and trigger T cell responses.

To investigate whether splenocytes from naïve animals can stimulate primary T cells, we generated T cell cultures from A/J mice immunized with SERCA2a 971-990. We evaluated T cell responses to splenocytes pulsed with or without SERCA2a 971-990 or MCC 82-103 (control) at a 1:1 ratio (Fig. 3A). We noted that the T cells co-cultured




Fig. 1. Characterization of hybridoma cells. (A) Antigen-specificity. Clone 80 hybridoma cells were stained with two sets of dextrans (IE^k/SERCA2a 971-990 and MCC 82-103; and IA^k/SERCA2a 971-990 and RNase 43-56) for 2 h, followed by anti-CD3, anti-CD4, and 7-AAD. After washing, cells were acquired by flow cytometry and the dext⁺ cells were determined in the live (7-AAD⁻) CD3⁺CD4⁺ subset. (B) Cytokine analysis. Hybridoma cells were stimulated with a cocktail of PMA/Ionomycin/GolgiStop for 5 h. After washing, cells were fixed and permeabilized to stain with the indicated cytokine antibodies and their corresponding controls. After acquiring by flow cytometry, percentages of cytokine⁺CD4⁺ T cells were determined. Representative data from 3 individual experiments are shown.




Fig. 2. Proliferative responses of SERCA2a 971-990 -specific hybridoma cells cultured with or without splenocytes from naïve mice and antigens. Clone 80 cells were cultured with splenocytes from A/J mice with or without SERCA2a 971-990 (specific) or MCC 82-103 (control) at different ratios for 24 h. After pulsing the cells with ³[H] thymidine for 16 h, proliferative responses were measured. Mean ± SEM values are shown (n = 3). *p ≤ 0.001, **p = 0.023.

with splenocytes alone responded significantly, and expectedly, SERCA2a 971-990, further increased the response, whereas responses obtained with T cells cultured with MCC 82-103 were comparable with those obtained with splenocytes alone (Fig. 3A). Further, we verified

the T cell responses to be MHC-restricted, as anti-IA^k and IE^k antibodies significantly blunted the proliferation of T cells co-cultured with APCs alone (Fig. 3B, left panel) or with SERCA2a 971-990, but not with control (Fig. 3B, right panel). The data suggest that the APCs from naïve

A





Fig. 3. Determination of proliferative responses of SERCA2a 971-990 -sensitized T cells to APCs loaded with or without peptides in the presence or absence of MHC antibodies. (A) Responses to APCs loaded with or without peptides. SERCA2a 971-990 -sensitized T cells were generated from A/J mice immunized with SERCA2a 971-990, and cells were cultured with SERCA2a 971-990 or MCC 82-103 in the presence or absence of splenocytes generated from A/J mice. Forty-eight hours later, cells were pulsed with ³[H] thymidine, and after 16 h, proliferative responses were measured. (B) Responses to APCs loaded with or without peptides in the presence or absence of MHC antibodies. Splenocytes from naïve mice were treated with Fc blocker and cultured with or without T cells in the absence (left panel) or presence of SERCA2a 971-990 (right panel) and anti-IA^k and anti-IE^k antibodies or rat IgG (control). Proliferative responses were measured as above after pulsing with ³[H] thymidine. Mean ± SEM values are shown (n = 3). *p ≤ 0.001.

B






Fig. 4. Cytokine responses in SERCA2a 971-990 -sensitized T cells exposed to splenocytes from naïve animals. SERCA2a 971-990 -sensitized T cells were cultured with splenocytes from naïve mice supplemented with or without SERCA2a 971-990 (specific) or MCC 82-103 (control) for two days where monensin was added during the last 12 h in each of the four different time points (18, 24, 36 and 48 h). At each point, cells were harvested and stained with anti-CD4, 7AAD. After fixation and permeabilization steps, cells were stained with three cocktails of antibodies, each representing IFN- γ -APC and IL-2-PE, IL-17A-APC and GM-CSF-PE and their corresponding isotype controls (rat IgG1-APC and rat IgG2a/IgG2b-PE). Cells were acquired by flow cytometry to analyze the percentages of cytokine-producing CD4 T cells within the live (7-AAD⁻) subset, where cells producing IFN- γ , IL-17A and their corresponding controls (all APC-conjugated) are shown in the Y axis and the cells producing IL-2, GM-CSF and their controls (all PE-conjugated) are shown in the X axis in the respective plots. Frequencies of cytokine producing cells are shown in the tables below each plot by adding the percentage values of the upper left and upper right quadrants for APC-conjugated cytokine antibodies (IFN- γ , IL-17A and their controls). Similarly, percentage values of the upper right and lower right quadrants were added for PE-conjugated cytokine antibodies (IL-2, GM-CSF and their controls). Representative data sets from 3 individual experiments are shown.

mice can induce proliferation of SERCA2a 971-990 -sensitized T cells through MHC class II pathway, thereby potentially modulating their functional responses.


3.2. SERCA2a 971-990 -sensitized T cells exposed to splenocytes from naïve mice produce inflammatory cytokines implicated in the development of autoimmune myocarditis

We had previously shown that the SERCA2a 971-990 -sensitized T cells generated from immunized animals produced mainly Th1 and Th17 cytokines that can transfer disease to naïve animals (Krishnan et al., 2018). We reasoned that the SERCA2a 971-990 -sensitized T cells exposed to APCs from naïve mice produce cytokines similar to those produced in response to SERCA2a 971-990 antigen-specifically. By intracellular cytokine staining, we analyzed the frequencies of cytokine-producing T cells in co-culture experiments, where we cultured SERCA2a 971-990 -sensitized T cells with splenocytes (APCs) from naïve animals in the absence or presence of SERCA2a 971-990 or MCC 82-103 (control). Cytokine analysis was performed at four time points (18, 24, 36 and 48 h) using antibody cocktails containing IFN- γ and IL-2, IL-17A and GM-CSF and their corresponding controls as described in the Methods (Section, 2.6). As shown in Fig. 4, T cells with no exposure to APCs, contain a low frequency of cytokine producing cells except IFN- γ , if any at 18 h (6.3 %). In contrast, cytokine-producing cells were absent in the cultures containing APCs-alone. However, upon exposure to APCs, SERCA2a 971-990 -sensitized T cells were found to produce the inflammatory cytokines implicated in the induction of various autoimmune diseases including myocarditis (Krishnan et al., 2018; Cihakova et al., 2008; Geppert and Lipsky, 1985; Bracamonte-Baran

and Čiháková, 2017; Baldeviano et al., 2010). These include mainly, IFN- γ (63.5 %) and GM-CSF (6.0 %), and, to a lesser extent, IL-17A (1.6 %) and IL-2 (1.2 %) with their peaks reaching at 24 h. As expected, cytokine responses were also upregulated and reached peaks at 18 or 24 h in the presence of SERCA2a 971-990 (IFN- γ , 69.3 %; GM-CSF, 20.5 %; IL-17A, 9.1 %; and IL-2, 4.2 %) (Fig. 4). Likewise, cytokine responses in cultures exposed to control (MCC 82-103) were also comparable to the T cells co-cultured with APCs alone during the same period (IFN- γ , 58.0 %; GM-CSF, 5.0 %; IL-17A, 1.6 %; and IL-2, 1.0 %). These data indicate that the SERCA2a 971-990 -sensitized T cells responded to the antigen displayed by the APCs and produce inflammatory cytokines that may have a pathogenic role in the development of autoimmune myocarditis.

3.3. Heart and liver APCs and ECs differentially modulate SERCA2a 971-990 -sensitized T cell responses

First, we asked whether APCs derived from non-lymphoid organs can also present SERCA2a 971-990 by isolating MNCs from the hearts and livers of naïve A/J mice. We obtained MNCs from groups of A/J mice, and, after staining with anti-CD45 and 7-AAD, we sorted CD45⁺CD3⁻ viable (7-AAD⁻) cells by flow cytometry. We noted that the CD45⁺ cells contained cells in the order of B cells (CD19⁺), followed by macrophages (CD11b⁺) and dendritic cells (CD11c⁺) and the MNCs also expressed the MHC class II molecule IE^k (Supplementary Fig. 1A). Next, we cultured both heart and liver MNCs with SERCA2a 971-990 -reactive T cells as described above. The data revealed enhanced proliferative responses when T cells were cultured with either MNCs alone or with peptides (SERCA2a 971-990 or MCC 82-103). Responses obtained with MCC 82-103 were comparable to those for



cells cultured with MNCs alone (Supplementary Fig. 1B), and these patterns were very similar to those noted with splenocytes described above. Collectively, the data suggest that APCs present in both lymphoid and non-lymphoid organs can display SERCA2a 971-990 and stimulate antigen-primed T cells. Of note, SERCA exists in multiple isoforms (Lompré, 1998; Brandl et al., 1987; Wuytack et al., 1995). While, SERCA2a is preferentially expressed in cardiomyocytes, vascular smooth muscle cells and ECs, SERCA2b can be expressed in APCs like macrophages (Li et al., 2004) that also has an equivalent epitope of SERCA2a 971-990. Thus, it is possible that epitope from multiple isoforms can potentially stimulate SERCA2a 971-990-sensitized T cells. It is also possible that the APCs in the target organs like heart can acquire self-antigens from other SERCA2a-expressing cells through phagocytosis of dead cells or autophagy that migrate to the spleens (Ansari et al., 1991; Larsen et al., 1990).

Next, since SERCA2a is known to be expressed in non-professional APCs, we tested a hypothesis that these cells are not expected to stimulate T cells. To this end, we purified CD31⁺ ECs by magnetic separation from hearts of naive A/J mice as previously described (Lim et al., 2003; Chou et al., 2015; van Beijnum et al., 2008). After verifying the purity of ECs to be more than 90 % (Fig. 5A), we used them in co-culture experiments using SERCA2a 971-990-reactive T cells in a proliferation assay. The data revealed that the T cell responses in cultures involving heart ECs and T cells were blunted (Fig. 5B, left panel). However, upon supplementation with SERCA2a 971-990, the T cell responses were further blunted, but not in the cultures exposed to MCC 82-103 (control) (Fig. 5B, left panel). We then tested whether liver ECs

Fig. 5. Determination of proliferative responses of SERCA2a 971-990-sensitized T cells cultured with or without ECs. (A) CD31 staining. Heparin was administered to groups of naive A/J mice, and after 10 min, animals were euthanized to collect hearts and livers. Organs were washed with ice-cold DPBS, and tissue homogenates were used to purify ECs as described in the Methods section. Cells were plated in culture dishes pre-coated with gelatin, and after 5–7 days they were examined for CD31 expression by flow cytometry. (B) Proliferative response. SERCA2a 971-990-sensitized T cells were cultured for 2 days with heart and liver ECs supplemented with or without SERCA2a 971-990 (specific) or MCC 82-103 (control). Cells were pulsed with ³[H] thymidine, and 16 h later proliferative responses were measured. Mean ± SEM values representing 3 individual experiments with 5–6 animals in each are shown. *p ≤ 0.001, **p = 0.023.

follow a pattern similar to heart ECs by sorting the ECs from liver (Fig. 5A, right panel). In co-culture experiments, ECs stimulated the SERCA2a 971-990-primed T cells (Fig. 5B, right panel), and the patterns of T cell responses were similar to those obtained with splenocytes as described above. The finding that the basal proliferative responses obtained with heart ECs were high (Fig. 5B), but reduced upon exposure to T cells, suggests that the heart ECs might have been suppressed by T cells leading to their death. But why, then, exposure to SERCA2a 971-990, but not MCC 82-103 led to reduced proliferative responses? Similarly, why such a pattern was lacking with the liver ECs? One possibility is that the ECs in various organs can behave differently, and their phenotypes can differ in the expression of MHC and co-stimulatory molecules (Lim et al., 2003; Pober et al., 2017).

To address the above questions, we examined the expression of MHC class II (IA^k and IE^k), MHC class I (H-2D^d, K^d and L^d), and co-stimulatory (CD80, CD86, and CD40) molecules in both heart and liver ECs by flow cytometry. These analyses revealed that heart ECs lacked the expression of both IA^k and IE^k, CD86, and CD40 and showed only low levels of CD80 (4.0 %) (Fig. 6, top left panel). Similar analysis of liver ECs showed the expression of IE^k, but not IA^k, whereas other molecules (CD80, CD86 and CD40) were detected in the range of ~60 to 90 % (Fig. 6, top right panel). Heart and liver ECs, however, expressed all three MHC class I molecules as expected (data not shown). We then exposed the heart and liver ECs to IFN-γ, a cytokine known to promote the expression of MHC class II and co-stimulatory molecules, to determine its effects on ECs (Linke and Male, 1994; Dore-Duffy et al., 1996; Knolle et al., 1999; Marelli-Berg et al., 2000; Ma and Pober,




Fig. 6. Effect of IFN- γ on ECs. Naive A/J mice were used to prepare ECs from hearts and livers as described in the Methods section, and cells were treated with or without IFN- γ for 3 days. After trypsinization, cells were washed and stained with the indicated markers and 7-AAD. Cells were acquired by flow cytometry to analyze the expression of indicated markers with the live (7-AAD⁻) CD31⁺ subset.

1998; Scott et al., 2018; Geppert and Lipsky, 1985). Expectedly, heart ECs treated with IFN- γ revealed upregulation of both MHC class II molecules (IA^k, 49.0 % and IE^k, 21.0 %), as well as CD80 (97.0 %) and CD40 (77.0 %), but expression of CD86 remained unaltered (Fig. 6, bottom left panel). Similarly, expression of the MHC class II molecule IA^k in liver ECs, which was lacking prior to exposure to IFN- γ , could now be detected with IFN- γ treatment (Fig. 6, bottom right panel). Overall, the data suggest that lack of expression of MHC class II molecules in heart ECs correlates with the absence of proliferative responses of T cells, since MHC class II molecules are needed for antigen presentation. Conversely, since liver ECs express at least one of the two MHC class II molecules (IE^k), they are able to present SERCA2a 971-990 to the responding T cells. This may be the reason the proliferative responses obtained in cultures containing liver ECs and T cells merely tended to be greater (Fig. 5B, right panel). While these explanations can be logically construed, it was difficult to explain why the proliferative responses of ECs were blunted in the heart. One possibility is that the SERCA2a 971-990 may contain antigenic determinants for CD8 T cells, and if so, they may potentially induce EC death. Identification of CD8 T cell determinants, if any may provide new insights as to their significance.

In summary, by creating T cell hybridomas for SERCA2a 971-990, we addressed a hypothesis that naïve APCs can spontaneously induce T cell responses antigen specifically. This hypothesis was tested using APCs generated from both lymphoid and non-lymphoid organs, and demonstrate that the responding T cells can produce pro-inflammatory cytokines. Of note, presentation of MHC/peptide complexes has been previously described for other cardiac antigens like cardiac myosin heavy chain- α (Smith and Allen, 1992), but the pathologic significance was not tested. Likewise, constitutive presentation of such complexes has also been reported for self-antigens of the central nervous system such as myelin basic protein (MBP) 1-44, MBP 83-102, PLP 139-151 and PLP 178-191 (Voskuhl, 1998). Furthermore, we had demonstrated that A/J mice although contain SERCA2a 971-990 -reactive T cells in their naïve periphery (Krishnan et al., 2018), they remain healthy (Krishnan et al., 2018). It is currently unknown whether enhanced production of inflammatory cytokines (in particular, IL-17A) that occurs in the immunized animals is a missing factor (Krishnan et al., 2018). Alternatively, antigen-specific T cells in the naïve animals may have a capacity to induce disease, but they could be potentially suppressed by regulatory T cells. We have not addressed these possibilities. While our data have provided an explanation as to how SERCA2a 971-990 -reactive T cells can induce myocarditis (Krishnan et al., 2018), the data also revealed new insights into the possibility that SERCA2a 971-990 -reactive T cells can potentially induce EC death. This may be an important mechanism because EC dysfunction has been proposed as an important predisposing factor for induction of vascular inflammation and coronary heart diseases like atherosclerosis (Tefamariam and DeFelice, 2007; Steyers and Miller, 2014; Gimbrone and García-Cardeña, 2016; Tousoulis et al., 2008).

Author statement

RA, BY, CM, XS and JR: Conceived and designed the experiments; RA, BY, SA, NL, and SJ: performed the experiments; JJR, analyzed the data; and RA, XS and JR: wrote the paper.

Funding

This work was supported by the National Institutes of Health [HL114669]. JJR is supported by funds from the Nebraska Research Initiative to the Bioinformatics Core Research Facility at the Center for Biotechnology.

Declaration of Competing Interest

All other authors declare no competing interests.

Acknowledgments

RA, BY, CM, XS and JR: Conceived and designed the experiments; RA, BY, SA, NL, and SJ: performed the experiments; JJR, analyzed the data; and RA, XS and JR: wrote the paper.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.imbio.2019.12.005>.

References

- Alanio, C., Lemaitre, F., Law, H.K., Hasan, M., Albert, M.L., 2010. Enumeration of human antigen-specific naïve CD8⁺ T cells reveals conserved precursor frequencies. *Blood* 115 (18), 3718–3725. <https://doi.org/10.1182/blood-2009-10-251124>. PubMed PMID: 20200354.
- Ansari, A., Wang, Y., Danner, D., Gravanis, M., Mayne, A., Neckelmann, N., et al., 1991. Abnormal expression of histocompatibility and mitochondrial antigens by cardiac tissue from patients with myocarditis and dilated cardiomyopathy. *Am. J. Pathol.* 139 (2), 337.
- Ashwell, J., Jenkins, M., Schwartz, R., 1988. Effect of gamma radiation on resting B lymphocytes. II. Functional characterization of the antigen-presentation defect. *J. Immunol.* 141 (8), 2536–2544.
- Baldeviano, G.C., Barin, J.G., Talor, M.V., Srinivasan, S., Bedja, D., Zheng, D., et al., 2010. Interleukin-17A is dispensable for myocarditis but essential for the progression to dilated cardiomyopathy. *Circ. Res.* 106 (10), 1646–1655.
- Basavalingappa, R.H., Massilamany, C., Krishnan, B., Gangaplara, A., Kang, G., Khalilzad-Sharghi, V., et al., 2016. Identification of an epitope from Adenine nucleotide translocator 1 that induces inflammation in heart in A/J mice. *Am. J. Pathol.* 186 (12), 3160–3175. <https://doi.org/10.1016/j.ajpath.2016.08.005>. PubMed PMID: 27876151; PubMed Central PMCID: PMC5225289.
- Basavalingappa, R.H., Massilamany, C., Krishnan, B., Gangaplara, A., Rajasekaran, R.A., Afzal, M.Z., et al., 2017. beta1-adrenergic receptor contains multiple IA(k) and IE(k) binding epitopes that induce t cell responses with varying degrees of autoimmune myocarditis in A/J mice. *Front. Immunol.* 8 (1567). <https://doi.org/10.3389/fimmu.2017.01567>. PubMed PMID: 29209317; PubMed Central PMCID: PMC5701947.
- Bracamonte-Baran, W., Čiháková, D., 2017. Cardiac autoimmunity: myocarditis. *The Immunology of Cardiovascular Homeostasis and Pathology*. Springer, pp. 187–221.

- Brandl, C., Martin, D., MacLennan, D., 1987. Adult forms of the Ca²⁺ ATPase of sarcoplasmic reticulum. Expression in developing skeletal muscle. *J. Biol. Chem.* 262 (8), 3768–3774.
- Bridgett, M., Cetkovic-Cvrlje, M., O'Rourke, R., Shi, Y., Narayanswami, S., Lambert, J., et al., 1998. Differential protection in two transgenic lines of NOD/Lt mice hyper-expressing the autoantigen GAD65 in pancreatic beta-cells. *Diabetes* 47 (12), 1848–1856.
- Bright, J.J., Xin, Z., Sriram, S., 1999. Superantigens augment antigen-specific Th1 responses by inducing IL-12 production in macrophages. *J. Leukoc. Biol.* 65 (5), 665–670.
- Cheng, M., Anderson, M.S., 2018. Thymic tolerance as a key brake on autoimmunity. *Nat. Immunol.* 1.
- Chou, C.H., Lai, S.L., Ho, C.M., Lin, W.H., Chen, C.N., Lee, P.H., et al., 2015. Lysophosphatidic acid alters the expression profiles of angiogenic factors, cytokines, and chemokines in mouse liver sinusoidal endothelial cells. *PLoS One* 10 (3), e0122060 Epub 2015/03/31. doi: 10.1371/journal.pone.0122060. PubMed PMID: 25822713; PubMed Central PMCID: PMC4379007.
- Cihakova, D., Barin, J.G., Afanasyeva, M., Kimura, M., Fairweather, D., Berg, M., et al., 2008. Interleukin-13 protects against experimental autoimmune myocarditis by regulating macrophage differentiation. *Am. J. Pathol.* 172 (5), 1195–1208.
- Daneker, G.W., Lund, S.A., Caughman, S.W., Swerlick, R.A., Fischer, A.H., Staley, C.A., et al., 1998. Culture and characterization of sinusoidal endothelial cells isolated from human liver. *In Vitro Cell. Dev. Biol. Anim.* 34 (5), 370–377 PubMed PMID: 9639099.
- Donermeyer, D.L., Beisel, K.W., Allen, P.M., Smith, S.C., 1995. Myocarditis-inducing epitope of myosin binds constitutively and stably to I-Ak on antigen-presenting cells in the heart. *J. Exp. Med.* 182 (5), 1291–1300 PubMed PMID: 7595200; PubMed Central PMCID: PMC2192215.
- Dore-Duffy, P., Balabanov, R., Rafols, J., Swanborg, R., 1996. Recovery phase of acute experimental autoimmune encephalomyelitis in rats corresponds to development of endothelial cell unresponsiveness to interferon gamma activation. *J. Neurosci. Res.* 44 (3), 223–234.
- French, M.B., Allison, J., Cram, D.S., Thomas, H.E., Dempsey-Collier, M., Silva, A., et al., 1997. Transgenic expression of mouse proinsulin II prevents diabetes in nonobese diabetic mice. *Diabetes* 46 (1), 34–39.
- Gangaplara, A., Massilamany, C., Brown, D.M., Delhom, G., Pattnaik, A.K., Chapman, N., et al., 2012. Coxsackievirus B3 infection leads to the generation of cardiac myosin heavy chain- α -reactive CD4 T cells in A/J mice. *Clin. Immunol.* 144 (3), 237–249. <https://doi.org/10.1016/j.clim.2012.07.003>. PubMed PMID: 22854287.
- Gangaplara, A., Massilamany, C., Steffen, D., Reddy, J., 2013. Mimicry epitope from *Ehrlichia canis* for interphotoreceptor retinoid-binding protein 201–216 prevents autoimmune uveoretinitis by acting as altered peptide ligand. *J. Neuroimmunol.* 263 (1–2), 98–107.
- Geppert, T., Lipsky, P., 1985. Antigen presentation by interferon-gamma-treated endothelial cells and fibroblasts: differential ability to function as antigen-presenting cells despite comparable Ia expression. *J. Immunol.* 135 (6), 3750–3762.
- Gimbrone Jr., M.A., García-Cardena, G., 2016. Endothelial cell dysfunction and the pathobiology of atherosclerosis. *Circ. Res.* 118 (4), 620–636.
- Goodnow, C.C., Sprent, J., Fazekas de St Groth, B., Vinuesa, C.G., 2005. Cellular and genetic mechanisms of self tolerance and autoimmunity. *Nature* 435 (7042), 590–597 Epub 2005/06/03. doi: 10.1038/nature03724. PubMed PMID: 15931211.
- Graf, M., Auch-Schwell, W., Graf, K., Terbeek, D., Hertel, H., Unkelbach, M., et al., 1994. Isolation and characterization of macrovascular and microvascular endothelial cells from human hearts. *Am. J. Physiol.* 267 (6 Pt 2), H2138–48. <https://doi.org/10.1152/ajpheart.1994.267.6.H2138>. PubMed PMID: 7810713.
- Graf, M., Graf, K., Auch-Schwell, W., Terbeek, D., Hertel, H., Fleck, E., 1993. Cultivation and characterization of micro- and macrovascular endothelial cells from the human heart. *Eur. Heart J.* 14 (Suppl 1), 74–81 PubMed PMID: 8293783.
- Huynh, H.K., Oger, J., Dorovinizis, K., 1995. Interferon- β downregulates interferon- γ -induced class II MHC molecule expression and morphological changes in primary cultures of human brain microvessel endothelial cells. *J. Neuroimmunol.* 60 (1–2), 63–73.
- Knolle, P.A., Germann, T., Treichel, U., Uhrig, A., Schmitt, E., Hegenbarth, S., et al., 1999. Endotoxin down-regulates T cell activation by antigen-presenting liver sinusoidal endothelial cells. *J. Immunol.* 162 (3), 1401–1407.
- Krishnan, B., Massilamany, C., Basavalingappa, R.H., Gangaplara, A., Kang, G., Li, Q., et al., 2017. Branched chain alpha-ketoacid dehydrogenase kinase 111-130, a T cell epitope that induces both autoimmune myocarditis and hepatitis in A/J mice. *Immun. Inflamm. Dis.* 5 (4), 421–434. <https://doi.org/10.1002/iid3.177>. PubMed PMID: 28597552; PubMed Central PMCID: PMC5691315.
- Krishnan, B., Massilamany, C., Basavalingappa, R.H., Gangaplara, A., Rajasekaran, R.A., Afzal, M.Z., et al., 2018. Epitope mapping of SERCA2a identifies an antigenic determinant that induces mainly atrial myocarditis in A/J mice. *J. Immunol.* 200 (2), 523–537. <https://doi.org/10.4049/jimmunol.1701090>. PubMed PMID: 29229678; PubMed Central PMCID: PMC5760440.
- Krishnan, B., Massilamany, C., Basavalingappa, R.H., Rajasekaran, R.A., Kuszynski, C., Switzer, B., et al., 2015. Versatility of using major histocompatibility complex class II dextramers for derivation and characterization of antigen-specific, autoreactive T cell hybridomas. *J. Immunol. Methods* 426, 86–94. <https://doi.org/10.1016/j.jim.2015.08.007>. PubMed PMID: 26268454; PubMed Central PMCID: PMC4651793.
- Larsen, C.P., Morris, P.J., Austyn, J.M., 1990. Migration of dendritic leukocytes from cardiac allografts into host spleens. A novel pathway for initiation of rejection. *J. Exp. Med.* 171 (1), 307–314. <https://doi.org/10.1084/jem.171.1.307>. PubMed PMID: 2404081; PubMed Central PMCID: PMC4651793.
- Li, Y., Ge, M., Ciani, L., Kuriakose, G., Westover, E.J., Dura, M., et al., 2004. Enrichment of endoplasmic reticulum with cholesterol inhibits sarcoplasmic-endoplasmic reticulum calcium ATPase-2b activity in parallel with increased order of membrane lipids implications for depletion of endoplasmic reticulum calcium stores and apoptosis in cholesterol-loaded macrophages. *J. Biol. Chem.* 279 (35), 37030–37039.
- Lim, Y.C., Garcia-Cardena, G., Allport, J.R., Zervoglos, M., Connolly, A.J., Gimbrone Jr., M.A., et al., 2003. Heterogeneity of endothelial cells from different organ sites in T-cell subset recruitment. *Am. J. Pathol.* 162 (5), 1591–1601. [https://doi.org/10.1016/S0002-9440\(10\)64293-9](https://doi.org/10.1016/S0002-9440(10)64293-9). PubMed PMID: 12707043; PubMed Central PMCID: PMC1851196.
- Linke, A., Male, D., 1994. Strain-specific variation in constitutive and inducible expression of MHC class II, class I and ICAM-1 on rat cerebral endothelium. *Immunology* 82 (1), 88.
- Lohse, A.W., Dinkelmann, M., Kimmig, M., Herkel, J., Meyer zum Buschenfelde, K.H., 1996. Estimation of the frequency of self-reactive T cells in health and inflammatory diseases by limiting dilution analysis and single cell cloning. *J. Autoimmun.* 9 (5), 667–675. <https://doi.org/10.1006/jaut.1996.0087>. PubMed PMID: 8933283.
- Lompré, A.-M., 1998. The sarco (endo) plasmic reticulum Ca²⁺-ATPases in the cardiovascular system during growth and proliferation. *Trends Cardiovasc. Med.* 8 (2), 75–82.
- Ma, W., Pober, J.S., 1998. Human endothelial cells effectively costimulate cytokine production by, but not differentiation of, naive CD4⁺ T cells. *J. Immunol.* 161 (5), 2158–2167.
- Marelli-Berg, F.M., Scott, D., Bartok, I., Peek, E., Dyson, J., Lechler, R.I., 2000. Activated murine endothelial cells have reduced immunogenicity for CD8⁺ T cells: a mechanism of immunoregulation? *J. Immunol.* 165 (8), 4182–4189.
- Massilamany, C., Gangaplara, A., Basavalingappa, R.H., Rajasekaran, R.A., Khalilzad-Sharghi, V., Han, Z., et al., 2016. Localization of CD8 T cell epitope within cardiac myosin heavy chain- α 334–352 that induces autoimmune myocarditis in A/J mice. *Int. J. Cardiol.* 202, 311–321.
- Massilamany, C., Krishnan, B., Reddy, J., 2015. Major Histocompatibility Complex Class II Dextramers: New Tools for the Detection of antigen-Specific, CD4 T Cells in Basic and Clinical Research. *Scand. J. Immunol.* 82 (5), 399–408.
- Massilamany, C., Gangaplara, A., Steffen, D., Reddy, J., 2011a. Identification of novel mimicry epitopes for cardiac myosin heavy chain- α that induce autoimmune myocarditis in A/J mice. *Cell. Immunol.* 271 (2), 438–449.
- Massilamany, C., Upadhyaya, B., Gangaplara, A., Kuszynski, C., Reddy, J., 2011b. Detection of autoreactive CD4 T cells using major histocompatibility complex class II dextramers. *BMC Immunol.* 12 (40). <https://doi.org/10.1186/1471-2172-12-40>. PubMed PMID: 21767394; PubMed Central PMCID: PMC3151213.
- Massilamany, C., Gangaplara, A., Chapman, N., Rose, N., Reddy, J., 2011c. Detection of cardiac myosin heavy chain- α -specific CD4 cells by using MHC class II/IA k tetramers in A/J mice. *J. Immunol. Methods* 372 (1), 107–118.
- McDouall, R.M., Yacoub, M., Rose, M.L., 1996. Isolation, culture, and characterisation of MHC class II-positive microvascular endothelial cells from the human heart. *Microvasc. Res.* 51 (2), 137–152. <https://doi.org/10.1006/mvvr.1996.0016>. PubMed PMID: 8778570.
- Merrick, A., Errington, F., Milward, K., O'Donnell, D., Harrington, K., Bateman, A., et al., 2005. Immunosuppressive effects of radiation on human dendritic cells: reduced IL-12 production on activation and impairment of naive T-cell priming. *Br. J. Cancer* 92 (8), 1450.
- Pober, J.S., Merola, J., Liu, R., Manes, T.D., 2017. Antigen presentation by vascular cells. *Front. Immunol.* 8, 1907.
- Prasad, S.A., Gregerson, D.S., 1997. Antigen presentation in uveitis. *Eye* 11 (2), 176.
- Reddy, J., Bettelli, E., Nicholson, L., Waldner, H., Jang, M.-H., Wucherpfennig, K.W., et al., 2003. Detection of autoreactive myelin proteolipid protein 139–151-specific T cells by using MHC II (IAs) tetramers. *J. Immunol.* 170 (2), 870–877.
- Rock, K.L., Rothstein, L., Gamble, S., 1990. Generation of class I MHC-restricted T-T hybridomas. *J. Immunol.* 145 (3), 804–811 PubMed PMID: 2115542.
- Sakaguchi, S., Sakaguchi, N., Asano, M., Itoh, M., Toda, M., 1995. Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor α -chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J. Immunol.* 155 (3), 1151–1164 PubMed PMID: 7636184.
- Scott, N.A., Zhao, Y., Krishnamurthy, B., Maninger, S.I., Kay, T.W., Thomas, H.E., 2018. IFN γ -Induced MHC Class II Expression On Islet Endothelial Cells Is An Early Marker Of Insulinitis But Is Not Required For Diabetogenic CD4⁺ T Cell Migration. *Front. Immunol.* 9.
- Skowronski, J., Jolicoeur, C., Alpert, S., Hanahan, D., 1990. Determinants of the B-cell response against a transgenic autoantigen. *Proc. Natl. Acad. Sci.* 87 (19), 7487–7491.
- Smith, S.C., Allen, P.M., 1992. Expression of myosin-class II major histocompatibility complexes in the normal myocardium occurs before induction of autoimmune myocarditis. *Proc. Natl. Acad. Sci. U. S. A.* 89 (19), 9131–9135. <https://doi.org/10.1073/pnas.89.19.9131>. PubMed PMID: 1409614; PubMed Central PMCID: PMC50079.
- Steinert, E.M., Schwartz, R.H., Singh, N.J., 2012. At low precursor frequencies, the T-cell response to chronic self-antigen results in anergy without deletion. *Eur. J. Immunol.* 42 (11), 2875–2880. <https://doi.org/10.1002/eji.201242518>. PubMed PMID: 22806568; PubMed Central PMCID: PMC3725465.
- Steyers, C., Miller, F., 2014. Endothelial dysfunction in chronic inflammatory diseases. *Int. J. Mol. Sci.* 15 (7), 11324–11349.
- Tamura, N., Ishii, N., Nakazawa, M., Nagoya, M., Yoshinari, M., Amano, T., et al., 1996. Requirement of CD80 and CD86 molecules for antigen presentation by eosinophils. *Scand. J. Immunol.* 44 (3), 229–238.
- Tesfamariam, B., DeFelicis, A.F., 2007. Endothelial injury in the initiation and progression of vascular disorders. *Vascul. Pharmacol.* 46 (4), 229–237. <https://doi.org/10.1016/j.vph.2006.11.005>. PubMed PMID: 17218160.
- Theofilopoulos, A.N., Kono, D.H., Baccala, R., 2017. The multiple pathways to autoimmunity. *Nat. Immunol.* 18 (7), 716.
- Tousoulis, D., Charakida, M., Stefanadis, C., 2008. Endothelial function and inflammation

- in coronary artery disease. *Postgrad. Med. J.* 84 (993), 368–371.
- van Beijnum, J.R., Rousch, M., Castermans, K., van der Linden, E., Griffioen, A.W., 2008. Isolation of endothelial cells from fresh tissues. *Nat. Protoc.* 3 (6), 1085–1091 Epub 2008/06/13. PubMed PMID: 18546599.
- Voskuhl, R.R., 1998. Myelin protein expression in lymphoid tissues: implications for peripheral tolerance. *Immunol. Rev.* 164 (1), 81–92.
- Vrisekoop, N., Artusa, P., Monteiro, J.P., Mandl, J.N., 2017. Weakly self-reactive T-cell clones can homeostatically expand when present at low numbers. *Eur. J. Immunol.* 47 (1), 68–73. <https://doi.org/10.1002/eji.201646540>. PubMed PMID: 27792294.
- Wuytack, F., Dode, L., Baba-Aissa, F., Raeymaekers, L., 1995. The SERCA3-type of organellar Ca²⁺ pumps. *Biosci. Rep.* 15 (5), 299–306.